

Chemical Engineering 412

Introductory Nuclear Engineering

Lecture 15

Industrial Applications

Exam II Review



Spiritual Thought


“Repentance is one of the first principles of the gospel. Forgiveness is a mark of divinity. There is hope for you. Your lives are ahead, and they can be filled with happiness, even though the past may have been marred by sin. This is a work of saving and assisting people with their problems. This is the purpose of the gospel.

This is the time, this is the very hour, to repent of any evil in the past, to ask for forgiveness, to stand a little taller and then to go forward with confidence and faith.”

President Gordon B. Hinkley



Roadmap



Economics

America derives substantial economic and employment benefits from the use of radiation and radioactive materials:



**\$330.7 billion annually
in total industrial
sales**

4,000,000 jobs




**\$60 billion in tax
revenues to local, state &
federal governments**



Economics

Nuclear energy's direct and indirect economic impacts in the US:

442,000 jobs



**\$90 billion in total
sales of goods &
services**





**\$17.8 billion in local, state
& federal tax revenues**

Destination

Once they are produced, they are packaged and shipped safely to users throughout the United States; users are:



Universities



Industries


Hospitals



Laboratories

Scientific Research


The FDA requires that all new drugs be tested for safety and effectiveness; more than 80% are tested with radioactive materials




Radioactive materials are also used in biomedical research, metabolic studies, genetic engineering and environmental protection studies

Scientific Research

Archaeologists use ^{14}C to date artifacts containing plant or animal material




Criminal investigators use radiation to examine evidence



Museums rely on radioactive materials to verify authenticity of art objects and paintings


Industrial Uses

Automobile industry makes use of isotopes to test the quality of steel in cars




Aircraft manufacturers use radiation to check for flaws in jet engines

Mining & petroleum companies use isotopes to locate and quantify geological mineral deposits



Industrial Uses

Oil gas & mining companies use isotopes to map geological contours (using test wells) and mine bores and to determine presence of hydrocarbons



Pipeline companies utilize radioactive isotopes to look for defects in welds

Construction crews use radioactive materials to gauge soil moisture content and asphalt density



Agricultural Uses

Hardier and more disease resistant crops (peanuts, tomatoes, onions, rice, soybeans, barley) have been developed using radioactive materials in agricultural research




Nutritional value, baking and melting qualities of some crops and cooking times have been improved using isotopes

Radioactive materials pinpoint where illnesses strike animals to breed disease-resistant livestock




Agricultural Uses

Radioactive materials show how plants absorb fertilizer; this helps researchers figure where and how much to apply to crops for maximum yield




Isotopes help farmers and scientists control pests; e.g., California has used radiation sterilization since the mid-70s to control Mediterranean fruit fly infestations

Consumer Products & Services



**96 US nuclear power plants
provide ~20% of electricity**

**Smoke detectors installed in
~90% of America's homes rely
on 1-2 μCi of ^{241}Am to monitor
for smoke to signal a fire**



**Computer disks retain data
better when treated with radiation**

Consumer Products & Services



Non-stick pans are treated with radiation to retain the coating

Photocopiers and plastic manufacturers use small amounts of radiation to eliminate static and prevent jamming



Cosmetics, hair products and contact lens solutions are sterilized with radiation to remove irritants and allergens

Consumer Products & Services

Radioactive materials are used to sterilize medical bandages and implements as well as foodstuffs to kill pathogens



1930s Fiestaware contains uranium in the ceramic glazes

To maximize light output, some lantern mantles contain radioactive thorium nitrate



The Large Hadron Collider


LHC is located at CERN
CERN is located near **Geneva**
Part of CERN is in France

The LHC collides protons
Center of Mass $E=14\text{ TeV}$ $\sim 7\times$ Fermilab
Very high luminosity $\sim 100\times$ Fermilab

Goal: discover Higgs+SUSY+???




The Large Hadron Collider



The Large Hadron Collider

Overall view of the LHC experiments.




ES40 - V10/09/97


The Large Hadron Collider

LHC PROJECT


SURFACE BUILDINGS



Above Ground



Below Ground



The Large Hadron Collider

Magnetic field at 7 TeV: 8.33 Tesla

Operating temperature: 1.9 K

Number of magnets: ~9300

Number of main dipoles: 1232



Number of quadrupoles: ~858

Number of correcting magnets: ~6208

Number of RF cavities: 8 per beam;


Field strength at top energy ≈ 5.5 MV/m

Power consumption: ~120 MW




How Do We Get 7 TeV Protons?

LINAC → PSB → PS → SPS → LHC




$\sim 10^{11}$ protons/beam



LHC: Large Hadron Collider
 SPS: Super Proton Synchrotron
 AD: Antiproton Decelerator
 ISOLDE: Isotope Separator OnLine Device
 PSB: Proton Synchrotron Booster
 PS: Proton Synchrotron
 LINAC: LINear ACcelerator
 LEIR: Low Energy Ion Ring
 CNGS: Cern Neutrinos to Gran Sasso

CERN Accelerators (not to scale)



Radolf LEY, PS Division, CERN, 02.09.96
 Revised and adapted by Antonella Del Rosso, ITT Div.,
 in collaboration with B. Desforges, SL Div., and
 D. Manghji, PS Div. CERN, 23.05.01

Chapter 8

- Detector Types
 - Dead times, interaction rates, performance, paralyzable, etc.
 - Examples & Diagrams, etc.
 - Fundamental operation principles
- Detection and Operation Modes
- Spectroscopy
- Efficiency
- Related equipment (PMTs, SCPHA, MCA)



Chapter 9

- Know Big picture of Radiation Doses
- Know various measurements, units conversion from one to another
 - KERMA, exposure, Absorbed Dose, etc.
 - Know how to correlate to biological impacts
- Calculation of dose
- Hazards of Radiation (Table usage)
- Exposure limits – amounts, history, etc.
- Perspective on radiation effects
- Acute and latent effects/symptoms
- Does model – Linear, threshold, hormesis



Chapter 13

- Beneficial Uses of Radiation +Applications
 - Specific isotopes and production
- Advantages/Disadvantages of radioactive
- Uses of Tracers (calculate amount needed)
- Uses of “Materials affecting Radiation”
- Uses of “Radiation affecting Materials”
- Particle Accelerators
- Economics and Widespread applications



Chapter 14

- Medical Uses of Radiation
 - Diagnostic vs. therapeutic
- X-Rays
- Mammography & Densitometry
- CT Scan
- SPECT
- PET
- MRI



Example I

Mercury pollution in water is of concern since fish often concentrate this element in their tissues. To measure such mercury contamination by neutron activation analysis, a fish sample is irradiated in a reactor in a thermal neutron flux $\phi = 1.5 \times 10^{12} \text{ cm}^{-2}\text{s}^{-1}$. The stable mercury isotope ^{196}Hg has a neutron absorption cross section for thermal neutrons of 3.2 kb. The resulting ^{197}Hg has a half life of 2.67 d and can be detected in a fish sample at a minimum activity of 15 Bq. What irradiation time is needed to detect mercury contamination at a level of 30 ng/g (30 ppm) in a 10-g fish sample?

$$A_{\text{cum}} = \text{in} - \text{out} + \text{gen} - \text{cons}$$

$$\frac{dN^{197}}{dt} = R_a - A \Rightarrow N \lambda^{197}$$

$$\frac{dN^{197}}{dt} = R_a - N \lambda^{197}$$

$$R_a = \phi \cdot \sigma_a^{196} \cdot N^{196}$$

$$N = \frac{P N_A}{A} = \frac{\text{atoms}}{\text{cm}^3}$$

$$M_m = \frac{30 \text{ ng/g}}{\text{g fish}}$$

300 ng Hg $\Rightarrow N = \frac{m}{M} \cdot N_A = 1.383 \times 10^{12}$ atoms Hg

$\frac{dN}{dt} = -\lambda N$ (with λ atoms) $\Rightarrow N = N_0 e^{-\lambda t}$ (with λ atoms)

$\frac{dN}{dt} + N\lambda = C$

$N_H + N_P = N$

$N_P = C$

$N' + N = C$

$N_H' = \frac{dN_H}{dt}$

$N_H' + N_H = 0$

$N_H = C_1 e^{-\lambda t}$

$N_P' + N_P = C$

$N_P = C - N_H$

\hat{R}_a

↓

↓

↓

$N(t) = \frac{\hat{R}_a}{\lambda} (1 - e^{-\lambda t})$

$A(t) = \hat{R}_a (1 - e^{-\lambda t})$

Example II

- If potatoes receive gamma-ray doses between 60 Gy and 150 Gy, premature sprouting is inhibited. Such irradiation can be done in an irradiator with a large ^{60}Co source. Assume all the gamma rays are absorbed in the potatoes. What is the minimum activity of ^{60}Co needed in an irradiator to deliver such a dose of 250kGy to 100,000 kg of potatoes in 8 hours?

- From appendix D, $E_\gamma = 2.5 \text{ MeV}$

$$E_{abs} = \frac{2.5 \times 10^5 \text{ J/kg} \times 10^5 \text{ kg}}{8 \times 3600 \text{ s} \times 1.602 \times 10^{-13} \text{ J/MeV}} = 5.42 \times 10^{18} \text{ MeV/s.}$$

Dose-mass = $E / 2.5 \text{ MeV} = \text{doses}$

- $A = 2.17 \text{E}18 \text{ Bq}$

$$N = \frac{m \cdot N_a}{A} \quad m = \frac{N \cdot A}{N_a}$$

Example III

- A Lithium-6/Hydrogen-3 sample is bombarded with neutrons (in a flux of $4.5\text{E}11$ neutrons/cm²/s) for 15 hours to determine how much is present in a given sample. The cross section of Lithium 6 is 943 b.
b. If the final number of Lithium 7 atoms is $N = 4.3\text{E}18$ atoms, what was the original number of ${}^6\text{Li}$ atoms?

