

# Chemical Engineering 412

## *Introductory Nuclear Engineering*

### Lecture 8

### Radiation Interactions with Matter

### Exam Review




# Spiritual Thought

“If it was right when you prayed about it and trusted it and lived for it, it is right now. Don’t give up when the pressure mounts. Certainly don’t give in to that being who is bent on the destruction of your happiness. Face your doubts. Master your fears. “Cast not away therefore your confidence.” Stay the course and see the beauty of life unfold for you.

Jeffery R. Holland



# Roadmap



# Objectives

- Be able to calculate probabilities of interaction and radiation field intensities
- Understand both **linear interaction coefficients** and **cross-sections**
- Be able to calculate or find  $\mu$ ,  $\sigma$ , and  $\Sigma$
- Be able to read, understand and take values from cross section libraries: plots or tables
- **Know how to calculate reaction rates!!!**



# Conceptual Interpretations

- The linear attenuation coefficient can be thought of in three ways:
  - Probability that a particle interacts in a differential length of material (does not assume constant  $\mu_i$ )
  - Inverse of the mean free path of a particle (assumes constant  $\mu_i$ ).
  - Related to distance at which half of particles have interacted ( $x_{1/2, i} = \frac{\ln 2}{\mu_i}$ ) (assumes constant  $\mu_i$ )
- Analogous to decay constants
  - Decay probabilities
  - Average lifetimes
  - Half lives.



# Flux and Reaction Rate

- If  $\sigma$  is probability of one particle interacting with one nucleus
- And  $\Sigma$  is the probability of one particle interacting with many nuclei
- How do we evaluate many particles with many nuclei?
- FLUX- Essentially particle density per time
- Reaction Rate (number of reactions per volume per time)

$$\hat{R}_i = \phi \sum_i \sigma_i = \phi N \sigma_i = \phi \sigma_i \frac{\rho N_a}{A}$$



# Example

- What is the power generation in a  $1\text{cm}^3$  section of  $\text{U}^{235}$  fuel, assuming a thermal neutron flux of  $1 \times 10^{22}$  neutrons/ $\text{cm}^2\text{-s}$ ?

From book:  $\sigma_f = 587\text{b}$   $N = \frac{\rho N_A}{M}$   $M = 235\text{g/mol}$   
 Appendix C  $\rho = 19.1\text{g/cm}^3$   $N = 4.89 \times 10^{22} \frac{\text{atoms}}{\text{cm}^3}$

$$E_f \approx 200\text{MeV}$$

$$\Sigma_f = \sigma \cdot N =$$

$$\text{Power} = \Sigma_f \cdot \Phi \cdot E_f = \underline{9.206\text{MW}}$$



# Mass Interaction Coefficient

- Photons – mass interaction coefficient
  - Interaction coefficient (macroscopic) divided by density
  - which depends only weakly on the properties of the medium (for photons)

$$\frac{\mu_i}{\rho} = \frac{\sigma_i N}{\rho} = \frac{N_a}{A} \sigma_i$$

- Homogeneous mixture properties can be determined from

$$\mu_i = \sum_j \mu_{i,j} = \sum_j N_j \sigma_{i,j} \quad \frac{\mu_i}{\rho} = \sum_j w_j \left( \frac{\mu_i}{\rho} \right)_j$$





# Cross Section Trends



- Most Isotopes
  - Cross sections rise as neutron energy decreases.
  - Resonance regions with narrow and rapidly varying interactions that eventually are not resolvable
- Light isotopes ( $A < 25$ )
- Heavy isotopes ( $A > 150$ )
- Intermediate




# Al Total Neutron Cross Section



# Fe total neutron cross section




# Lead Total Neutron Cross Section




# Fission Cross Sections $^{235}\text{U}$

Uranium 235 Fission Cross Section MT = 18


note scale




# Fission Cross Sections $^{235}\text{U}$




# Fission Cross Sections $^{235}\text{U}$




# Fission Cross Sections $^{235}\text{U}$




# Fission Cross Sections $^{235}\text{U}$



# Cross section over entire range




# Fission Cross Section of $^{238}\text{U}$



threshold energy (> resonance region energy)



# Fissionable Cross Sections



# Chemical Engineering 412

## *Introductory Nuclear Engineering*

### Exam 1 Review



# Chapter 1 - Fundamentals

- Nuclear units
- Elementary particles/particle physics
- Isotopic nomenclature
- Atomic weight/number density
- Chart of nuclides
- Mass energy equivalency



# Chapter 2 – Quantum Mechanics

- Special Relativity – time, length, mass changes
- Relativistic mass/momentum/energy relations
- Particle-wave duality
- Schrödinger's wave equation
- Heisenberg's uncertainty principle



# Chapter 3 – Nuclear Models

- Nuclear energy states
- Liquid Drop Model
- Nuclear mass equation
- Shell Model
- Nuclear stability
- Binding energy/mass excess
- Modern Nucleus concepts



# Chapter 4 – Nuclear Energetics

- Terminology
- Mass defect/BE
- Nuclear reactions
- Conserved quantities for various situations (not all the same!)
- **\*\*\*\*Q-Value\*\*\*\* (know how to calculate for ALL reactions)**
  - Know how to deal with charge
  - Know how to deal with excited nuclei
  - Know how to deal with electrons/binding energy of electrons



# Chapter 5 – Nuclear Decay

- Conservations
- Decay mechanisms – distinguishing features, Q values, energy/momentum balances
- \*\*\*Energy Diagrams\*\*\*
- Alpha/Beta particle energy distribution
- Decay Constant
- Half-Life
- Activity



# Chapter 5 – Nuclear Decay (cont)

- Parallel/Series Decay Routes
- Decay Chains
- Solutions to decay chain equations  
Secular Equilibrium
- Radionuclides in nature
- Carbon 14 dating
- Other isotopic dating methods
- Three component decays

Isobars and most stable masses



# Chapter 6 – Binary Nuclear Reactions

28

- Definitions
- Types of binary reactions
- Reaction Mechanisms
- Kinematics (scattering example)
- Threshold Energy
- Neutron Reactions
- Neutron Scattering/slowing
- Neutron Energy Spectrums



Lethargy

# Chapter 6 – Binary Nuclear Reactions (cont.)

- Neutron capture vs. slowing
- Fission reactions
- Emitted/recoverable fission energy
- Critical energies for fission
- Fertile vs. fissile vs. fissionable
- Fission product distribution
- Prompt vs. delayed neutrons
- Fission steps/timeline



# Chapter 7 – Radiation Interactions with matter

30

- Linear Interaction Coefficient (micro vs. macro)
- Cross section (micro vs. macro)
- Attenuation in Material
- Derivation of material interaction
- Buildup factor
- Mass Attenuation Coefficient
- Energy dependence of cross sections
- Cross section Trends



# Chapter 7 – Radiation Interactions with matter (cont.)

31

- Cross Section of mixture
- Total intensity/flux
- Neutron flux
- Time/space/position dependence of flux
- Fluence
- Uncollided flux transmission
- Thermal vs. fast neutrons
- Photon Interactions – types, trends, energies, cross sections
- Charged Particle interactions
- Stopping Power (collision vs. radiative)
- Range



# Example 1

- The Radionuclide  $^{41}\text{Ar}$  decays by  $\beta^-$  emission to an excited level of  $^{41}\text{K}$  that is 1.293 MeV above the ground state. What is the maximum kinetic energy of the emitted  $\beta^-$  particle?
- What makes this the maximum energy?



# Solution

## Q Equation

$$Q_{\beta^-} = \{M(^{41}_{18}\text{Ar}) - [M(^{41}_{19}\text{K}) + E^*/c^2]\}c^2$$

$$[40.9645008 - 40.9618259] * 931.5 - 1.293 \text{ MeV}$$

$$= 1.199 \text{ MeV}$$

B) Because an antineutrino is also released, which carries away some energy – this maximum is when the antineutrino has zero energy



# Example 2

Assume a fuel rod has a diameter of 1 cm and a length of 5 m. Assuming an enrichment of 5%  $^{235}\text{U}$  and a thermal flux of  $2 \times 10^{13}$  neutrons, what is the reaction rate in the fuel rod for:

- a) scattering?
- b) fission?



## Example 2 key

- $\sigma_{s235} = 13.7\text{b}$ ,  $\sigma_{f235} = 587\text{b}$
- $\sigma_{s238} = 9.47\text{b}$ ,  $\sigma_{f238} = 0\text{b}$
- $N_{235} = 2.447\text{E}21 \text{ cm}^{-3}$ ,  $N_{238} = 4.591\text{E}22 \text{ cm}^{-3}$
- $V_f = 392.7 \text{ cm}^3$
- $R_s = 3.678\text{E}15\text{s}^{-1}$
- $R_f = 1.128\text{E}16\text{s}^{-1}$



## Example 3

- What is the probability of producing  $^{91}\text{Br}$  in a fission reaction?
- Use fission product mass distribution chart:
- $\sim 8.5\%$



# Example 4

- What is the amount of thermal neutrons that are absorbed in water per  $\text{cm}^3$  over 1 hour in a fission reactor if the thermal flux is  $2.2 \cdot 10^{16}$  neutrons/ $\text{cm}^2/\text{s}$ ? ( $\Sigma_a = 0.0197 \text{ cm}^{-1}$ )

- $1.56 \cdot 10^{18}$  absorptions per  $\text{cm}^3$

$$\Phi \cdot t = \bar{\Phi}$$

$$\hat{R}_a = \Phi \cdot \Sigma_a \text{ or } \bar{\Phi} \cdot \Sigma_a$$



# Example 5

